



Identification of *Elizabethkingia meningoseptica* from American bullfrog (*Rana catesbeiana*) farmed in Sabah, Malaysia using PCR method and future management of outbreak

Julian Ransangan^{1*}, Noramini Zainuri¹, Tamrin Mohammad Lal¹, Boniface Jintoni² and Vui Sheong Chung²

¹Microbiology and Fish Disease Laboratory, Borneo Marine Research Institute, Universiti Malaysia Sabah, 88400, Jalan UMS, Kota Kinabalu, Sabah, Malaysia.

²Fisheries Department Sabah, Wisma Pertanian, Jalan Tasik, 88628 Kota Kinabalu, Sabah, Malaysia.
E-mail: liandra@ums.edu.my

Received 28 April 2012; Received in revised form 17 July 2012; Accepted 19 July 2012

ABSTRACT

Aims: High demand for frog meat in Malaysia especially the American bullfrog (*Rana catesbeiana*) has promoted intensive farming of the animal. However, the farming of American bullfrog is restricted by the occurrence of diseases. This study reports the first isolation of *Elizabethkingia meningoseptica* from specimens of American bullfrog that suffer from cataract and 'red-leg' syndrome.

Methodology and Result: The pathogen was isolated from eyes and internal organs (liver, kidney and spleen) of the diseased bullfrog specimens. All the bacterial isolates were subjected to phenotypic characterization and antibiotic susceptibility assay, and further identified by using the 16S rDNA sequencing analysis. We designed two pair of specific PCR primers (22-25 mers) which are complimentary to the β -lactamase gene in the reference strain of *E. meningoseptica* ATCC49470. The result showed all the bacterial isolates shared similar phenotypic characters and antibiotic susceptibility. BLAST analysis of the 16S rDNA sequences indicated that the bacterial isolates had very high sequence homology (100%) with *E. meningospetica* ATCC49470 and *E. meningoseptica* isolates from mosquito. The two PCR primers were very specific to *E. meningoseptica* isolates of this study.

Conclusion, significance and impact of study: This is the first isolation and characterization of bacterial pathogen, *E. meningoseptica* in cultured American bullfrog (*Rana catesbeina*) that suffered from eye cataract and 'red-leg' syndrome in Sabah, Malaysia. It is suspected that one of the possible transmission routes of the bacterial pathogen could be via mosquito bites. The findings suggest that there is urgent requirement for standard guideline of good farming practice to be adopted in frog farms throughout the country. Such a guideline can help in minimizing economic losses, preventing transmission of the zoonotic bacterial pathogen to farm workers, and sustaining the industry in Malaysia and upgrading frog meat quality for international market.

Keywords: American bullfrog, *Elizabethkingia meningoseptica*, cataract, red-leg syndrome, Malaysia

INTRODUCTION

American bullfrog, *Rana catesbeiana*, is not native to Malaysia but it was introduced in the country for aquaculture purposes. The farming of this anuran species has gained popularity because of its excellent attributes of adaptability to various tropical environmental conditions and relatively large size, with rich muscle mass. In Malaysia, the American bullfrog is farmed commercially to satisfy both local and international markets. Initially, there were only 12 bullfrog farms operating in Malaysia with the annual production of 80 tons of anuran meat (Kechick, 1995). After 14 years of introduction, the frog meat industry in Malaysia has undergone much development whereby the current meat production is estimated at 100

tons per month (Lee *et al.*, 2009). However, in the recent years the production of bullfrog in some farms in Malaysia is limited by the high mortality due to diseases. The most frequently occurring diseases are the 'red-leg' syndrome and cataract. These diseases affect mainly the adult frogs. The cataract is characterized by opaque eye lens, lethargic behaviour and loss of appetite. The frog with red-leg syndrome seemed to have limited hopping ability with no appetite for food. These two diseases can sometimes concurrently occur in the same individual frog. The diseases caused mortality within few days to weeks after the onset. Previous study on various tissues of bullfrog with redleg syndrome showed presence of variety bacterial pathogens which include *Aeromonas hydrophila*, *Elizabethkingia* (*Chryseobacterium*) *meningoseptica*,

Streptococcus innie, *Edwardsiella tarda*, *Citrobacter frundii* and *Pseudomonas* spp. (Mauel *et al.*, 2002). Contrary to the previous finding, this study reports the isolation of single bacterial pathogen, *E. meningoseptica* from bullfrog with redleg and cataract. In addition, this study also describes the specific PCR method for detection of the bacterial pathogen.

MATERIALS AND METHODS

Bacterial isolation and preservation

Bacteria were isolated from internal organs (liver, spleen and kidney) and eyes of the diseased frogs. Briefly, the frogs were aseptically dissected using sterile surgical tools to expose internal organs including spleen, kidney, heart and liver, and eyes. Sterile inoculating loop was aseptically swabbed on each organ, streaked on tryptic soy agar (TSA, Difco, USA) plates that were supplemented with 1.5% (w/v) sodium chloride and incubated at 28 °C for 48 h. Subsequently, the bacteria were serially sub-cultured on TSA plates to obtain single pure colony. Finally, 5 well characterized bacterial isolates were preserved at -86 °C according to the method described by Floodgate and Hayes (1961).

Phenotypic characterization

The 5 bacterial isolates were subjected to various biochemical tests according to the method described by Ransangan and Mustafa (2009). Tests included Gram staining, motility test, oxidative-fermentative test, catalase test, oxidase test, acid and gas production from sugars, citrate utilization, urease test, methyl-red reaction, Voges Proskauer, indole production, phenylalanine test, β -galactosidase test, lysine decarboxylase and arginine dehydrolase. The bacterial isolates were also grown at four different temperatures (10 °C, 28 °C, 37 °C and 40 °C) in four concentrations (0, 2, 4 and 6%) of NaCl (w/v).

Antibiotic susceptibility test

The bacterial isolates were then subjected to antibiotic susceptibility assay. They were first grown on TSA plates for 24 h at 28 °C. Each bacterial isolate was suspended in sterile phosphate buffered saline (PBS) (pH 7.2) and diluted to a turbidity equivalent to a MacFarland No. 0.5 standard solution. Then, 0.1 mL bacterial suspension was spread onto Mueller-Hinton agar (Difco) plate. Subsequently, antibiotic discs were aseptically placed onto the inoculated plates according to the method described by Dalsgaard *et al.* (1999). The antibiotic disks (Oxoid, Hampshire, England) used in this assay included ampicillin (10 μ g), chloramphenicol (30 μ g), ciprofloxacin (5 μ g), sulphamethoxazole (100 μ g), furazolidone (100 μ g), kanamycin (30 μ g), nalidixic acid (3 μ g), neomycin (10 μ g), nitrofurantoin (300 μ g), novobiocin (5 μ g), oxolinic acid (2 μ g), oxytetracycline (30 μ g), penicillin G (10 units), streptomycin (25 μ g), sulphonamide (300 μ g), tetracycline (10 μ g) and vancomycin (30 μ g). The plates were

incubated at 28 °C for 48 h and inhibition zones were scored according to the method described by Barry *et al.*, (1979).

DNA isolation

Genomic DNA from the bacterial isolates was extracted using the CTAB-DTAB method as described by Phillips and Simon (1995). First, each bacterium was inoculated in 5 mL sterile tryptic soy broth (TSB, Difco) and incubated overnight at 28 °C overnight following the method described by Kim and Jeong (2001). Subsequently, 1.0 mL of the bacterial suspension was transferred into microtube and centrifuged at 7,500 g for 5 min at 4 °C. After centrifugation, the supernatant was discarded, the bacterial pellet was re-suspended in 600 μ L DTAB solution [8% DTAB; 1.5 M NaCl; 100 mM Tris-HCl (pH 8.8) and 50 mM EDTA] and incubated at 75 °C for 5 min. The mixture was added with 700 μ L of chloroform and vortex for 20-30 sec before centrifugation at 13,400 g for 5 min. Later on, 450 μ L of the aqueous layer was transferred into the new sterile microtube, and added with 100 μ L CTAB solution (5% CTAB; 0.4 M NaCl) and 900 μ L sterile dH₂O. The mixture was incubated again at 75 °C for 5 min and centrifuged at 13,400 g for 10 min. Supernatant was discarded and DNA pellet dissolved in 150 μ L dH₂O and incubated further at 75 °C for 5 min before centrifugation at 13,400 g for 5 min. Once again the clear solution was transferred into new microtube and added with equal volume of 95% ethanol. The microtube was finger flicked several times and centrifuged again at 13,400 g for 10 min. The DNA pellet was washed with 75% ethanol and centrifuged at similar speed as described above. Finally, the DNA pellet was dissolved in 50 μ L 1X TE buffer and stored at -20 °C until use. The DNA concentration was determined using GeneQuant Pro RNA/DNA calculator (Pharmacia).

PCR amplification of 16S rRNA gene

PCR amplification of 16S rRNA gene was conducted against total genomic DNA extracted from all the 5 isolates using primers shown in Table 1. The forward and reverse primers were correspondent to nucleotide positions 3776045 to 3776026 and 3774654 to 3774678 of the 16S rRNA gene of *E. coli* ATCC 8739 (Figure 1), respectively. The PCR The PCR amplification was conducted in 25 μ L total reaction which consisted of 2.5 μ L of 10X i-Taq PCR buffer (iNtRON, Korea), 1.0 μ L of each (10 μ M) forward and reverse primers, 0.5 μ L i-Taq Polymerase (iNtRON), 2.0 μ L DNA template (0.307 μ g/ μ L l) and 18.0 μ L nuclease-free water. The amplification was carried out one cycle at 95 °C for 3 min followed by 30 cycles at 95 °C for 1 min, at 58 °C for 1 min and at 72 °C for 1 min, and final extension at 72 °C for 5 min. The PCR products were analyzed on 1.5% agarose gel electrophoresis and visualized under UV using the Alphaimager[®] Imaging System (Alpha Innotech Corporation).

Table 1: PCR primers used in this study.

Primer	Nucleotide Sequence (5'-3')	Target gene	Expected size (bp)
16SFJR	ATBNAGAGTTTGATCMTGGC	16S rRNA	1400
16SRJR	CAAGGCCCGGAACGTATTCAC	16S rRNA	
JREMF1	ATATTACGTAGGAACCTATGATTTG	β -lactamase	612
JREMR1	ATGGAGATCGAACTGACTTGCAT	β -lactamase	
JREMF2	ATGATTTGGCTTCTTACCTTATTG	β -lactamase	644
JREMR2	TATCCATAAACCAATTGCGGATT	β -lactamase	



Figure 1: Locations of PCR primers (16SFJR: nt3776045 to nt3776026; 16SRJR: nt3774654 to nt3774678) of 16S rRNA gene within the genome of *Escherichia coli* ATCC 8739 (CP000946).

DNA cloning and sequencing

PCR products (16S rDNA fragments) were purified using *AccuPrep*TM. PCR purification Kit (Bioneer Corporation, Seoul, Korea) according to the procedures described in the manufacturer's instruction manual. Two microlites (2.0 μ L) of the PCR product was cloned into pGEM[®]-T Easy (Promega, Madison, USA) cloning vector following the method described by the manufacturer. The plasmid was purified using PureLinkTM Quick Plasmid Miniprep Kit (InvitrogenTM, USA) following manufacturer's instruction. The plasmids were restricted using EcoR1 (New England Biolabs, USA) and analyzed on 1.5% agarose gel electrophoresis. Finally, 20 μ L of each purified plasmid harbouring correct fragment of the 16S rDNA was sequenced using M13 primers (Macrogen, DNA sequencing service, Seoul, Korea). Bacterial isolates were identified based on the result of BLAST analysis of the partial 16S rDNA sequences. The percentage identity

of 16S rDNA sequences of the 5 bacterial isolates against 16S rDNA sequences downloaded from the genbank was computed using the ClustalW (DNASTAR, Madison, United States). The construction of phylogenetic tree was achieved using the MegAlign program (DNASTAR) and the TREECON for Windows (Van de peer and De Wachter, 1994).

Specific PCR detection of *E. meningoseptica*

Two PCR primer pairs were designed based on β -lactamase gene sequences downloaded from genbank (DQ004496, GU188445, EF394442, EF394444, EF394445 and EF394446). The first primers pair was correspondent to the nucleotide positions, 132 to 256 and 744 to 722 of *E. meningoseptica* GOB-18 gene (DQ004496) and second primers pair was correspondent

to the nucleotide positions, 149 to 172 and 793 to 772 of *E. meningoseptica* GOB-18 gene (DQ004496) (Figure 2). The PCR amplification was carried out in 25 μ L total reaction which consisted of 12.5 μ L 10X i-Taq PCR buffer (iNtRON), 1.0 μ L of each primer (10 μ M), 0.3 μ L i-Taq polymerase (iNtRON), 2.0 μ L DNA (0.307 μ g/ μ L) and 8.2 μ L sterile Milli-Q water. The optimum PCR condition when using primers (JREMF1 and JREMR1) was as follow: initial DNA denaturation at 95 $^{\circ}$ C for 3 min followed by 30 cycles at 95 $^{\circ}$ C for 30 sec, 58 $^{\circ}$ C for 30 sec and 72 $^{\circ}$ C for 30 sec, and final extension at 72 $^{\circ}$ C for 5 min. Whereas the optimum PCR condition when using primers (JREMF2 and JREMR2) was as described above except for the annealing temperature set at 55 $^{\circ}$ C.

The specificity of the primers was evaluated against DNA from *Vibrio alginolyticus* (ATCC 17749), *V. parahaemolyticus* (ATCC 17802), *V. harveyi* (ATCC 35084), *V. anguillarum* (ATCC 19264), *Aeromonas salmonicida* subsp. *salmonicida* (ATCC33658), *A. hydrophila* (ATCC 7965), *A. caviae* (ATCC 15468), *Edwardsiella tarda* (ATCC 15947), *Yersinia ruckeri* (ATCC 29473), *Pseudomonas fluorescens* (ATCC 13525), *Pseudomonas aeruginosa* (ATCC 27853), *Proteus mirabilis* (ATCC 29245), *Escherichia coli* (ATCC 25922) and *E. meningoseptica* (ATCC 13253). The PCR products were analyzed on 1.5% agarose gel electrophoresis and visualized under UV using the Alphaimager[®] Imaging System.



Figure 2: Locations of β -lactamase PCR primers (JREMF1: nt132 to nt156, JREMF2: nt149 to nt172, JREMR1: nt744 to nt722 and JREMR2: nt793 to nt772) within the complete coding sequence of *Elizabethkingia meningoseptica* class B carbapenemase COB-18 gene (DQ004496).

RESULTS

Phenotypic characterization

All the 5 bacterial isolates exhibited similarity in the phenotypic features such as Gram staining negative, non-motile, and positive for oxidase and catalase, and negative for Voges-proskauer reaction. Although the bacteria were not able to produce indole, they utilized citrate and gelatine. They did not produce arginine dihydrolyse, lysine decarboxylase and urease but produced β -galactosidase and cellobiose but not from other sugars. Nevertheless, the gas was not produced from all the sugars tested. The bacteria grew at 28 $^{\circ}$ C and 37 $^{\circ}$ C but not at 10 $^{\circ}$ C and 40 $^{\circ}$ C, respectively. The bacteria were tolerant to NaCl concentrations up to 4% (w/v) but inhibited at 6% (w/v). However, the 5 bacterial isolates differed from the ATCC strains of *E. meningoseptica* in the acid production from cellobiose, D-fructose, maltose, lactose and manitol. Details of the phenotypic characteristics of the bacterial isolates are shown in Table 2.

Table 2: Phenotypic features of the *E. meningoseptica* isolates from American bullfrog, *Rana catesbeiana* farmed in Sabah, Malaysia.

Characteristics	Bacterial Isolates					<i>E. meningoseptica</i> ^a
	EKMS1	EKML1	EKMK1	EKMLE1	EKMRE1	
Gram staining	-	-	-	-	-	-
Shape	rod	rod	rod	rod	rod	rod
Oxidase test	+	+	+	+	+	+
Catalase test	+	+	+	+	+	nd
Voges-Proskauer	-	-	-	-	-	nd
Indole production	-	-	-	-	-	+
Citrate utilization	+	+	+	+	+	+
Gelatine	+	+	+	+	+	nd
Arginine dihydrolase	-	-	-	-	-	nd
Lysine decarboxylase	-	-	-	-	-	nd
Phenylalanine agar	-	-	-	-	-	nd
*ONPG	+	+	+	+	+	nd
Methyl-Red	-	-	-	-	-	nd
Urease test	-	-	-	-	-	-
Growth at 10 °C	-	-	-	-	-	nd
28 °C	+	+	+	+	+	nd
37 °C	+	+	+	+	+	+
40 °C	-	-	-	-	-	nd
Growth at 0 % NaCl	+	+	+	+	+	nd
2% NaCl	+	+	+	+	+	nd
4% NaCl	+	+	+	+	+	nd
6% NaCl	-	-	-	-	-	nd
O/F glucose	O	O	O	O	O	nd
Gas (acid) from glucose	- (+)	- (+)	- (+)	- (+)	- (+)	nd(+)
D-fructose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(+)
Cellobiose	- (+)	- (+)	- (+)	- (+)	- (+)	nd(-)
Mannose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)
Sorbitol	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)
Arabinose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(-)
Dextrose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)
Sucrose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)
Maltose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(+)
Mannitol	- (-)	- (-)	- (-)	- (-)	- (-)	nd(+)
Lactose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(+)
Salicin	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)
Raffinose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)
Galactose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)
Rhamnose	- (-)	- (-)	- (-)	- (-)	- (-)	nd(nd)

Bacterial identification

The PCR primers designed in this study successfully amplified partial fragment of 16S rDNA from all the bacterial isolates. Based on the BLAST analysis, it was found that all the bacterial isolates had high nucleotide sequence homology (98-100%) to the 16S rDNA

sequences belonging to *E. meningoseptica* strains (Table 3, Figure 3). On this basis, the 5 bacterial isolates described here are identified as those of *E. meningoseptica*. The partial 16S rDNA sequences of the 5 bacterial isolates were deposited in GenBank (<http://www.ncbi.nih.gov>) with the accession numbers as shown in Table 4.

Table 3: Percentage similarity (above diagonal) of 16S rDNA sequences of E. meningoseptica in reference to nt75-1350 of the 16S rDNA sequence (X80724) of Escherichia coli ATCC25922

Table with 48 columns and multiple rows showing percentage similarity values between various E. meningoseptica and Escherichia coli strains. The diagonal elements represent 100% similarity for each strain.




Figure 3: Phylogenetic relationship of the E. meningoseptica isolated from American bullfrog farmed in Sabah, Malaysia with other closely related bacteria. The tree was constructed based on alignment of 16S rDNA sequences using ClustalW method (DNASTAR Ver. 5.05) at positions corresponding to the nucleotides 75- 1350 of the 16S rDNA of Escherichia coli ATCC 25922 (X80724).

Antibiotic susceptibility assay

The 5 bacterial isolates exhibited strong susceptibility to ciprofloxacin, nalidixic acid, compound sulphonamides and trimethoprim. However, they were resistant to nitrofurantion, chloramphenicol, ampicilin, oxytetracycline, tetracycline, streptomycin and kanamycin. The inhibition zones exhibited by individual antibiotics against the bacterial isolates are given in Table 5.

Specific PCR for detection of *E. meningoseptica*

The primers against the β -lactamase gene of *E. meningoseptica* were specific to the bacteria. All the bacterial isolates were successfully amplified using both pairs of the PCR primers with expected sizes. In contrast, the primer pairs did not amplify any of the ATCC bacterial strains tested in this study. The results of PCR amplification using the primer pair 1 and pair 2 are shown in Figure 4 and Figure 5, respectively.

Table 4: List of 16S rDNA sequences used in study.

Bacterial Strain	Accession No.	Reference
<i>E. meningoseptica</i>	EF426431	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i> ATCC 49470	AJ704544	Kim <i>et al.</i> 2005
<i>E. meningoseptica</i>	EF426426	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426430	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426432	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426428	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426434	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426427	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426433	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426425	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	EF426429	Lindh <i>et al.</i> 2008
<i>E. meningoseptica</i>	FJ816028	Kajla <i>et al.</i> 2010
<i>E. meningoseptica</i>	FJ816020	Kajla <i>et al.</i> 2010
<i>E. meningoseptica</i> ATCC 13255	AJ704542	Kim <i>et al.</i> 2005
<i>C. meningosepticum</i>	AF207072	Bellais <i>et al.</i> 2000
<i>E. meningoseptica</i> ATCC 51720	AJ704545	Kim <i>et al.</i> 2005
<i>E. meningoseptica</i> ATCC 13254	AJ704541	Kim <i>et al.</i> 2005
<i>C. meningosepticum</i>	AF207071	Bellais <i>et al.</i> 2000
<i>C. meningosepticum</i>	AF207070	Bellais <i>et al.</i> 2000
<i>C. meningosepticum</i>	AF207078	Bellais <i>et al.</i> 2000
<i>C. meningosepticum</i>	AF207079	Bellais <i>et al.</i> 2000
<i>E. meningoseptica</i>	HM748601	Kim <i>et al.</i> 2011
<i>E. meningoseptica</i> ATCC 13253 ¹	AJ704540	Kim <i>et al.</i> 2005
<i>E. meningoseptica</i>	NR_042267	Kim <i>et al.</i> 2005
<i>E. meningoseptica</i>	FJ839441	Su and Ming, 2010
<i>C. meningosepticum</i>	AF207075	Bellais <i>et al.</i> 2000
<i>C. meningosepticum</i>	AF207077	Bellais <i>et al.</i> 2000
<i>C. meningosepticum</i>	AY468477	Bernardet <i>et al.</i> 2005
<i>C. meningosepticum</i>	AF207076	Bellais <i>et al.</i> 2000
<i>E. miricola</i>	NR_036862	Kim <i>et al.</i> 2005
<i>C. meningosepticum</i>	AF207073	Bellais <i>et al.</i> 2000
<i>E. miricola</i> ATCC 33958	AJ704543	Kim <i>et al.</i> 2005
<i>C. meningosepticum</i>	AF207074	Bellais <i>et al.</i> 2000
<i>E. meningoseptica</i>	EF204460	Shak�d <i>et al.</i> 2010
<i>C. isbilliense</i>	AM159184	unpublished
<i>C. hominis</i>	AM423083	Vaneechoutte <i>et al.</i> 2007
<i>Echerichia coli</i> ATCC 25922	X80724	Cilia <i>et al.</i> 1996
<i>E. meningoseptica</i>	GU180602	This study
<i>E. meningoseptica</i>	GU180603	This study
<i>E. meningoseptica</i>	GU180604	This study
<i>E. meningoseptica</i>	GU180605	This study
<i>E. meningoseptica</i>	GU180606	This study

Table 5: Inhibition zone (cm) recorded on different antibiotics against different isolates of *E. meningoseptica*

Bacteria	F	C	CIP	AMP	OA	OT	TE	S	NA	K	S3	W
EKME1	0	0.9	2.0	0	0.8	1.2	0	0	2.1	0	2.7	1.8
EKME2	0	1.2	2.1	0	1.2	1.2	0	0	2.0	0	2.7	1.7
EKML1	0	1.1	2.2	0	1.0	1.1	0	0	2.0	0	2.8	1.5
EKMK1	0	1.1	2.1	0	1.0	1.0	0	0	2.0	0	2.9	2.0
EKMS1	0	1.4	2.5	0	1.4	1.2	0	0	2.5	0	2.5	1.5




Figure 4: PCR amplification (612bp) using primers (JREMF1 and JREMR1) designed specifically for *E. meningoseptica*. Lanes M: 1 kb DNA Ladder (Promega); lane 1: *E. meningoseptica* ATCC 13253, lane 2: *E. meningoseptica* isolate EKMK1; lane 3: *E. meningoseptica* isolate EKML1; lane 4: *E. meningoseptica* isolate EKMLE1; lane 5: *E. meningoseptica* isolate EKMRE1; lane 6: *E. meningoseptica* isolate EKMS1; lane 7: *A. caviae* ATCC 15468; lane 8: *A. hydrophila* ATCC 7965; lane 9: *A. salmonicida* subsp. *salmonicida* ATCC 33658; lane 10: *Ed. tarda* ATCC 15947; lane 11: *Escherichia coli* ATCC 25922; lane 12: *Pr. mirabilis* ATCC 29245; lane 13: *P. aeruginosa* ATCC 27853; lane 14: *P. fluorescens* ATCC 13525; lane 15: *V. alginolyticus* ATCC17749; lane 16: *V. anguillarum* ATCC 19264; lane 17: *V. harveyi* ATCC 35084; lane 18: *V. parahaemolyticus* ATCC 17802; lane 19: *Y. ruckeri* ATCC 29473 and lane 20: Sterile double distilled water.



Figure 5: PCR amplification (644bp) using primers ((JREMF2 and JREMR2) designed specifically for *E. meningoseptica*. Lanes M: 1 kb DNA Ladder (Promega); lane 1: *E. meningoseptica* ATCC 13253, lane 2: *E. meningoseptica* isolate EKMK1; lane 3: *E. meningoseptica* isolate EKML1; lane 4: *E. meningoseptica* isolate EKMLE1; lane 5: *E. meningoseptica* isolate EKMRE1; lane 6: *E. meningoseptica* isolate EKMS1; lane 7: *A. caviae* ATCC 15468; lane 8: *A. hydrophila* ATCC 7965; lane 9: *A. salmonicida* subsp. *salmonicida* ATCC 33658; lane 10: *Ed. tarda* ATCC 15947; lane 11: *Escherichia coli* ATCC 25922; lane 12: *Pr. mirabilis* ATCC 29245; lane 13: *P. aeruginosa* ATCC 27853; lane 14: *P. fluorescens* ATCC 13525; lane 15: *V. alginolyticus* ATCC17749; lane 16: *V. anguillarum* ATCC 19264; lane 17: *V. harveyi* ATCC 35084; lane 18: *V. parahaemolyticus* ATCC 17802; lane 19: *Y. ruckeri* ATCC 29473 and lane 20: Sterile double distilled water.

DISCUSSION

E. meningoseptica was previously known as *Flavobacterium meningosepticum* or *Chryseobacterium meningosepticum* (Kim *et al.*, 2005). It is a Gram-negative and non-fermenting bacterium which is widely distributed in nature. It constitutes common bacterial flora in freshwater (Vancanneyt *et al.*, 1994). However, the bacterium has also been isolated from diseased turtles, frogs and fish (Green *et al.*, 1999; Bernardet, 2006; Mael *et al.*, 2003; Bernardet *et al.*, 2005). The bacterium was also recognized as an occasional but serious opportunistic pathogen to human, giving rise to meningitis, pneumonia, septic arthritis, endocarditis and conjunctivitis (Bernardet *et al.*, 2006).

Cataract is the most prevalent disease in farmed anurans and it quickly spreads within a relatively short period of time (Xie *et al.*, 2010). It is characterized by opaque eye lens, ascites in peritoneal cavity, lethargy and torticollis (Xie *et al.*, 2010). In the present study, we also observed eye opacity, sluggish behavior and ascites in peritoneal cavity of frogs with cataract and red-leg syndrome. In addition, the infected frogs had limited hopping ability and they were observed to suffer mortality from a few days to weeks after the onset of the disease. The bacterial isolation and 16S rDNA sequencing analysis revealed that all the bacterial isolates from eyes and internal organs of frogs belonged to *E. meningoseptica*. Although this bacterium has previously been isolated from farmed tiger frog (*R. tigrina rugulosa*) in China with cataract (Xie *et al.*, 2010) and in African clawed frog *Xenopus laevis* (Bernardet, 2006), this is the first report from Malaysia.

The farming of bullfrog in Malaysia is conducted either in earthen ponds or in concrete tanks. Broodstocks were first imported from Taiwan (Lee *et al.*, 2009). Ever since, they are propagated and maintained by the Fisheries Department of Sabah. Currently, the department maintains about 200-250 frog brooders in one of its aquaculture stations in Penampang. Frog larvae from this station are distributed to small-scale farms throughout Sabah as part of the government subsidy program. The cataract and red-leg syndrome occurred in frogs maintained in this station as well as in several private farms throughout Penampang district. The diseases were observed to affect adult frogs. It was found that the rapid spread of the disease among frogs could have been contributed by poor farming practices in most farms. This was apparent since diseased frogs are not isolated from clinically healthy animals. Furthermore, water quality in the culture tank was poor as indicated by smell of decaying organic matter. Uneaten foods are not removed but let to decay in the culture tanks. Similarly, several decaying dead frogs were also observed in the tanks. The workers who maintain the farms are not protected since they are handling the frogs by their bare hands. This could particularly be hazardous since *E. meningoseptica* has been reported as an opportunistic but serious human pathogen (Bernardet, 2006) especially those with respiratory problem (Weaver *et al.*, 2010).

E. meningoseptica can be contracted by the frogs from several sources including soils (Ahmad *et al.*, 2009), water (Vandamme *et al.*, 1994) and even mosquitoes (Lindh *et al.*, 2008; Rani *et al.* 2009). In the prevailing situation in Sabah, we understand that the bacterium may have been transmitted through mosquitoes. This explains the 100% nucleotide sequence homology of the five *E. meningoseptica* isolated in this study to 16S rRNA gene sequences of *E. meningoseptica* isolates (EF426426, EF426427, EF426430, EF426432 and EF426434) from mosquitoes, *Anopheles gambiae* (Lindh *et al.*, 2008). Although no bacterial isolation was done from mosquito specimens, the poorly maintained rearing water in the farms can be a perfect place for mosquito breeding. This is supported by the observation of many mosquito larvae in the rearing tanks in the farms where the disease outbreaks occurred.

The bacterium seemed to have developed resistance towards β -lactam antibiotics which include penicillin, nitrofurantion, ampicillin, tetracycline, streptomycin and kanamycin. Hence, the choice of antibiotics for treatment of cataract and red-leg syndrome is limited except for ciprofloxacin and nalidixic acid. However, it has been shown that the use of ciprofloxacin in African clawed frog (*X. laevis*) showed that there was an increase of the antibiotic concentration in the habitat several hours after the administration (Howard *et al.*, 2010).

The PCR primers targeting β -lactamase gene described in the present study can be potentially be developed as a DNA-based diagnostic kit for *E. meningoseptica*. However, verification and validation of the technique are still required before such a diagnostic kit can be developed. Despite the availability of detection kit, we strongly believe that good farming practices should be carried all the time in order to effectively prevent disease outbreak from occurring in the farms. These include improving of quality of rearing water, avoiding of stagnation, preventing exposure to mosquitoes, regular tank cleaning, and removal of any uneaten food and dead animals from the culture tanks. Need for effective protection should also be explained to farm workers because of the zoonotic potential of *E. meningoseptica*. With these programs in place, disease outbreaks can be prevented or at least significantly minimized to curtail economic losses and to sustain the frog meat industry in the country

CONCLUSION

The first isolation of *E. meningoseptica* from frog farms in Sabah may form the basis for an extensive epidemiological study of the pathogen to be carried out throughout frog farms in Malaysia. We strongly believe that there is urgent requirement of standard guideline for good farming practice to be adopted in frog farms throughout the country. Such a guideline can help in minimizing economic losses, preventing transmission of the zoonotic bacterial pathogens to farm workers, and

sustaining the industry in Malaysia as well as upgrading of frog meat quality for international market.

ACKNOWLEDGEMENT

We express our sincere thanks and appreciation to Prof. Dr. Ahmed Al-Harbi for his valuable suggestion on the experimental design especially the phenotypic characterization and to Prof. Saleem Mustafa for his critical review of the manuscript. The study was funded by the Ministry of Higher Education Malaysia (MOHE) under the grant no. FRG0188-ST-2009.

REFERENCES

- Ahmad, N., Johri, S., Abdin, M.Z. and Qazi, G.N. (2009). Molecular characterization of bacterial population in the forest soil of Kashmir, India. *World Journal of Microbiology and Biotechnology* **25**: 107-113.
- Barry, A.L., Coyle, M.B., Thornsberry, C., Gerlach, E.H. and Hawkinson, R.W. (1979). Methods of measuring zones of inhibition with the Bauer-Kirby disk susceptibility test. *Journal of Clinical Microbiology* **10**:885-889.
- Bellais, S., Aubert, D., Naas, T. and Nordmann, P. (2000). Molecular and Biochemical Heterogeneity of Class B Carbapenem-Hydrolyzing β -Lactamases in *Chryseobacterium meningosepticum*. *Antimicrobial Agents and Chemotherapy* **44**:1878 - 1886.
- Bernardet, J.F. (2006). Minutes of the Meetings of the International Committee on Systematics of Prokaryotes and Subcommittee on the taxonomy of *Flavobacterium* and *Cytophaga*-like bacteria. *International Journal of Systematic and Evolutionary Microbiology* **56**:2946-2951
- Bernardet, J.F., Bruun, B. and Hugo, C. (2006). The genera *Chryseobacterium* and *Elizabethkingia*. In: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.H. and Stackebrandt, E., editors. The prokaryotes: a Handbook on the Biology of Bacteria, 3rd edition, New York: Springer pp. 638-676.
- Bernardet, J.F., Vancanneyt, M., Matte-Tailliez, O., Grisez, L., Tailliez, P., Bizet, C., Nowakowski, M., Kerouault, B. and Swings, J. (2005). Polyphasic study of *Chryseobacterium* strains isolated from diseased aquatic animals. *Systematic and Applied Microbiology* **28**:640 – 660
- Cilia, V., Lafay, B. and Christen, R. (1996). Sequence heterogeneities among 16s Ribosomal RNA sequences, and their effect on phylogenetic analyses at the species level. *Molecular Biology and Evolution* **13**: 451 – 461.
- Dalsgaard, I., Høi, L., Siebeling, J. and Dalsgaard, A. (1999). Indole-positive *Vibrio vulnificus* isolated from disease outbreaks on a Danish eel farm. *Diseases of Aquatic Organisms* **35**: 187-194
- Floodgate, G.D. and Hayes, P.R. (1961). The preservation of marine bacteria. *Journal of Applied Bacteriology* **24**:87-93.
- Green, S.L., Bouley, D.M., Tolwani, R.J., Waggie, K.S., Lifland, B.D., Otto, G.M. and Ferrell, J.E. (1999). Identification and management of an outbreak of *Flavobacterium meningosepticum* infection in a colony of South African clawed frog (*Xenopus laevis*). *Journal of the American Veterinary Medical Association* **214**:1833-1838.
- Howard, A.M., Papich, M.G., Felt, S.A., Long, C.T., McKeon, G.P., Bond, E.S., Torreilles, S.L., Luong, R.H. and Green, S.L. (2010). The pharmacokinetics of enrofloxacin in adult African clawed frogs (*Xenopus laevis*). *Journal of the American Association for Laboratory Animal Science* **49**:800-804.
- Kajla, M.K., Andreeva, O., Gilbreath III, T.M. and Paskewitz, S.M. (2010). Characterization of expression, activity and role in antibacterial immunity of *Anopheles gambiae* lysozyme c-1. *Comparative Biochemistry and Physiology Part B* **155**:201- 209.
- Kechik, I.A. (1995). Aquaculture in Malaysia. In: Bagarinao TU, Flores EEC, editors. Towards sustainable aquaculture in Southeast Asia and Japan, Philippines: SEAFDEC p. 125-135.
- Kim, K.K., Kim, M.K., Lim, J.H., Park, H.Y. and Lee, S.T. (2005). Transfer of *Chryseobacterium meningosepticum* and *Chryseobacterium miricola* to *Elizabethkingia* gen. nov. as *Elizabethkingia meningoseptica* comb. nov. and *Elizabethkingia miricola* comb. nov. *International Journal of Systematic and Evolutionary Microbiology* **55**:1287-1293.
- Kim, M.S. and Jeong, H.D. (2001). Development of 16S rRNA targeted PCR methods for the detection and differentiation of *Vibrio vulnificus* in marine environment. *Aquaculture* **193**:199-211.
- Kim, S.J., Cho, C.H. and Whang, K.S. (2011). Characterization of protease produced by *Elizabethkingia meningoseptica* CS2-1 and optimization of cultured conditions for amino acid production. *Journal of Applied Biological Chemistry* **54**:135-142
- Lee, S.W., Musa, N., Wendy, W., Nadirah, M., Faizah, S.H. (2009). Occurrence of heavy metals and antibiotic resistance in bacteria from internal organs of American bullfrog (*Rana catesbeiana*) raised in Malaysia. *Journal of Venomous Animal and Toxins including Tropical Diseases* **15**: 353-358.
- Lindh, J.M., Borg-Karlson, A.K. and Faye, I. (2008). Transstadial and horizontal transfer of bacteria within a colony of *Anopheles gambiae* (Diptera:

- Culicidae) and oviposition response to bacteria-containing water. *Acta Tropica* **107**: 242-250.
- Mauel, M.J., Miller, D.L., Frazier, K.S. and Hines II, M.E. (2002).** Bacterial pathogens isolated from cultured bullfrogs (*Rana catesbeiana*). *Journal of Veterinary Diagnostic Investigation* **14**:431-433.
- Phillips, A.J. and Simon, C. (1995).** Simple, efficient, and nondestructive DNA extraction protocol for arthropods. *Annals of the Entomological Society of America* **88**:281-283.
- Rani, A., Sharma, A., Rajagopal, R., Adak, T. and Bhatnagar, R.K. (2009).** Bacterial diversity analysis of larvae and adult midgut microflora using culture-dependent and culture-independent methods in lab-reared and field-collected *Anopheles stephensi*-an Asian malarial vector. *BMC Microbiology* **9**: 96.
- Ransangan, J. and Mustafa, S. (2009).** Identification of *Vibrio harveyi* isolated from diseased Asian seabass *Lates calcarifer* by use of 16S ribosomal DNA sequencing. *Journal of Aquatic Animal Health* **21**:150-155.
- Shaked, T., Hantis-Zacharov, E. and Halperu, M. (2010).** *Epilithonimonas lactis* sp. nov., isolated from raw cow's milk. *International Journal of Systematic and Evolutionary Microbiology* **60**:675-679.
- Su, Z.J. and Ming, D.S. (2010).** Detection of blaTem-116, ant(3'')-Iof resistant related genes in a *Chryseobacterium meningosepticum* strain in sputa from a severe chronic hepatitis patient (abstract in English). *Chinese Journal of Nosocomiology* doi: cnki:sun:zhyy.0.2010-06-011.
- Van de Peer, Y. and De Wachter, R. (1994).** TREECON for Windows: a software package for the construction and drawing of evolutionary trees for the Microsoft Windows environment. *Computer Applications in Biosciences* **10**:569-570.
- Vancanneyt, M., Segers, P., Hauben, L., Hommez, J., Devriese, L.A., Hoste, B., Vandamme, P. and Kersters, K. (1994).** *Flavobacterium meningosepticum*, a pathogen in birds. *International Journal Systematic and Evolutionary Microbiology* **32**:2398-2403.
- Vandamme, P., Bernardet, J.F., Segers, P., Kersters, K. and Holmes, B. (1994).** New perspectives in the classification of the flavobacteria: description of *Chryseobacterium* gen. nov., *Bergeyella* gen. nov., and *Empedobacter* nom. rev. *International Journal of Systematic and Bacteriology* **44**: 846-849.
- Vanechoutte, M., Kämpfer, P., Baere, T.D., Avesani, V., Janssens, M. and Wauters, W. (2007).** *Chryseobacterium hominis* sp. nov., to accommodate clinical isolates biochemically similar to CDC groups II-h and II-c. *International Journal of Systematic and Evolutionary Microbiology* **57**: 2623 – 2628.
- Weaver, K.N., Jones, R.C., Albright, R., Thomas, Y., Zambrano, C.H., Costello, M.M., Havel, J., Price, J. and Gerber, S.I. (2010).** Acute emergence of *Elizabethkingia meningoseptica* infection among mechanically ventilated patients in a long-term acute care facility. *Infection Control and Hospital Epidemiology* **31**: 54-58.
- Xie, Z.Y., Zhou, Y.C., Wang, S.F., Mei, B., Xu, X.D., Wen, W.Y. and Feng, Y.Q. (2009).** First isolation and identification of *Elizabethkingia meningoseptica* from cultured tiger frog, *Rana tigrina rugulosa*. *Veterinary Microbiology* **138**:140-